

Power-law Behavior of Strings Scattered from Domain-wall at High Energies and Breakdown of their Linear Relations

Chuan-Tsung Chan^{*}

Department of Physics, Tung-Hai University, Taichung, Taiwan, R.O.C.

Jen-Chi Lee[†]

Department of Electrophysics, National Chiao-Tung University, Hsinchu, Taiwan, R.O.C.

Yi Yang[‡]

*Department of Electrophysics, National Chiao-Tung University and Physics Division,
National Center for Theoretical Sciences, Hsinchu, Taiwan, R.O.C.*

(Dated: February 2, 2008)

Abstract

In contrast to the common wisdom, we discover that, instead of the exponential fall-off of the form factors with Regge-pole structure, the high energy scattering amplitudes of string scattered from Domain-wall behave as power-law with Regge-pole structure. This is to be compared with the well-known power-law form factors without Regge-pole structure of the D-instanton scatterings. This discovery makes Domain-wall scatterings an unique example of a hybrid of string and field theory scatterings. The calculation is done for bosonic string scatterings of arbitrary massive string states from D24-brane. Moreover, we discover that the usual linear relations of high-energy string scattering amplitudes at each fixed mass level break down for the Domain-wall scatterings. This result gives a strong evidence that the existence of the infinite linear relations, or stringy symmetries, of high-energy string scattering amplitudes is responsible for the softer, exponential fall-off high energy string scatterings than the power-law field theory scatterings.

^{*}Electronic address: ctchan@thu.edu.tw

[†]Electronic address: jcclee@cc.nctu.edu.tw

[‡]Electronic address: yyang@phys.cts.nthu.edu.tw

In contrast to the local quantum field theory, string theory is remarkable for its ultra-violet finiteness. Being a consistent quantum theory of gravity with no free parameter, it is conceivable that an huge symmetry gets restored at high energies. Recently high energy, fixed angle behavior of string scattering amplitudes [1, 2, 3] was intensively reinvestigated for massive string states at arbitrary mass levels [4, 5, 6, 7, 8, 9, 10, 11]. The motivation was to uncover the fundamental hidden stringy spacetime symmetry conjectured by Gross in 1988. An important new ingredient of this approach is the zero-norm states (ZNS) [12, 13, 14] in the old covariant first quantized (OCFQ) string spectrum. One utilizes the decoupling of ZNS in the high energy limit to obtain infinite linear relations, or stringy symmetries, among string scattering amplitudes. Moreover, these linear relations can be used to fix the ratios among high energy scattering amplitudes of different string states at each fixed mass level algebraically. This explicitly shows that there is only one independent component of high-energy scattering amplitude at each mass level. On the other hand, a saddle-point method was developed to calculate the general formula of tree-level high-energy scattering amplitudes of four arbitrary string states to verify the ratios calculated above. This general formula expresses all high-energy string scattering amplitudes in terms of four tachyon scattering amplitude as conjectured by Gross [2].

In addition to the infinite linear relations, there are two other fundamental characteristics of high energy string scattering amplitudes, which make them very different from those of field theory scatterings. The first one is the exponential fall-off behavior of the form factors of string scatterings in the high-energy limit in contrast to the power-law behavior of point-particle field theory scatterings. It is the main result of this letter that the infinite linear relations, or stringy symmetries, among high-energy string scattering amplitudes of different string states discussed above are responsible for this softer exponential fall-off behavior of string scatterings than the power-law behavior of field theory scatterings. The second one is the Regge-pole structure [11] in the high energy string scattering amplitudes due to the infinite number of resonances in the string spectrum. In the scattering processes of string scattered from D-brane [15, 16], it was claimed that the exponential fall-off form factors with Regge-pole structure is a general feature of all Dp -brane scatterings except D-instanton, which exhibits power-law form factor without Regge-pole structure, and thus resembles a field theory amplitude instead.

In our recent work [17], we have shown that the linear relations stated above persist for the

string/D p -brane scatterings with $p \geq 0$. The linear relations for the D-particle scatterings were explicitly demonstrated. All the high energy string/D p -brane scattering amplitudes with $p \geq 0$ behave as exponential fall-off as was claimed in Ref [15]. In this letter, in contrast to the common wisdom, we discover that, instead of the exponential fall-off behavior of the form factors with Regge-pole structure, the high-energy scattering amplitudes of string scattered from D24-brane, or Domain-wall, behave as *power-law* with Regge-pole structure. This is to be compared with the well-known power-law form factors without Regge-pole structure of the D-instanton scatterings stated above. This discovery makes Domain-wall scatterings an unique example of a hybrid of string and field theory scatterings. Our calculation will be done for bosonic string scatterings of arbitrary massive string states from D24-brane. Moreover, we discover that the usual linear relations [17] of high energy string scattering amplitudes at each fixed mass level breaks down for the Domain-wall scatterings. This result gives a strong evidence that the existence of the infinite linear relations, or stringy symmetries, of high energy string scattering amplitudes is responsible for the softer, exponential fall-off high-energy string scatterings than the power-law field theory scatterings.

At a fixed mass level $M_{op}^2 = 2(n-1)$ of 26D open bosonic string theory, it was shown that [7, 8] a four-point function is at the leading order at high-energy limit only for states of the following form (we use the notation of [18])

$$|n, 2m, q\rangle \equiv (\alpha_{-1}^T)^{n-2m-2q}(\alpha_{-1}^L)^{2m}(\alpha_{-2}^L)^q |0, k\rangle. \quad (1)$$

where $n \geq 2m + 2q, m, q \geq 0$. Note that, in the high energy limit, the scattering process becomes a plane scattering (however, see the Domain-wall scattering in the end of this letter). The state in Eq.(1) is arbitrarily chosen to be the second vertex of the four-point function. The other three points can be any string states. We have defined the normalized polarization vectors of the second string state to be [4, 5]

$$e_P = \frac{1}{M_{op}}(E_2, k_2, 0) = \frac{k_2}{M_{op}}, \quad (2)$$

$$e_L = \frac{1}{M_{op}}(k_2, E_2, 0), \quad (3)$$

$$e_T = (0, 0, 1) \quad (4)$$

in the CM frame contained in the plane of scattering. By using the decoupling of two types of ZNS in the high energy limit, it was shown that there exists infinite linear relations among

string scattering amplitudes [7, 8]

$$\mathcal{T}^{(n,2m,q)} = \left(-\frac{1}{M_{op}}\right)^{2m+q} \left(\frac{1}{2}\right)^{m+q} (2m-1)!! \mathcal{T}^{(n,0,0)}. \quad (5)$$

Moreover, these linear relations can be used to fix the ratios among high energy scattering amplitudes of different string states at each fixed mass level algebraically. Eq.(5) explicitly shows that there is only one independent high-energy scattering amplitudes at each fixed mass level.

We now turn to the case of D-brane scatterings. In our recent work [17], we discover that scatterings of bosonic massive closed string states at arbitrary mass levels from D-brane can be expressed in terms of the generalized hypergeometric function ${}_3F_2$ with special arguments, which terminates to a finite sum and, as a result, the whole scattering amplitudes consistently reduce to the usual beta function. For the simple case of D-particle, we explicitly calculated high-energy scattering amplitude of an incoming closed string tensor state $(\alpha_{-1}^T)^{n-2q} (\alpha_{-2}^L)^q \otimes (\tilde{\alpha}_{-1}^T)^{n-2q'} (\tilde{\alpha}_{-2}^L)^{q'} |0\rangle$ with $M^2 \equiv \frac{M_{closed}^2}{2\alpha'_{closed}} = 2(n-1) = -k_2^2$ and an outgoing tachyon state to be [17]

$$\begin{aligned} A_{D-Par} &= A_{D-Par}^{(0 \rightarrow 1)} + A_{D-Par}^{(1 \rightarrow \infty)} \\ &= 2(-1)^{a_0} E^{2n} \left(-\frac{1}{2M}\right)^{q+q'} B\left(a_0+1, \frac{b_0+1}{2}\right) \\ &= 2(-1)^{a_0} E^{2n} \left(-\frac{1}{2M}\right)^{q+q'} \frac{\Gamma(a_0+1)\Gamma(\frac{b_0+1}{2})}{\Gamma(a_0+\frac{b_0}{2}+\frac{3}{2})}, \end{aligned} \quad (6)$$

where the high energy beta function $B(a_0+1, \frac{b_0+1}{2})$ is independent of $q+q'$, and a_0, b_0 will be defined in Eqs.(13) and (14). Since the calculation of decoupling of high energy ZNS without D-brane remains the same as the calculation of string scattered from D-brane, it is a remarkable result that the ratios $\left(-\frac{1}{2M}\right)^{q+q'}$ in Eq.(6) for different high energy scattering amplitudes at each fixed mass level is consistent with Eq.(5) for the scattering without D-brane as we found before [7]. One notes that the exponential fall-off behavior in energy E is hidden in the high energy beta function. Since the arguments of $\Gamma(a_0+1)$ and $\Gamma(a_0+\frac{b_0}{2}+\frac{3}{2})$ in Eq.(6) are negative in the high-energy limit, one needs to use the well known formula

$$\Gamma(x) = \frac{\pi}{\sin(\pi x) \Gamma(1-x)} \quad (7)$$

to calculate the large negative x expansion of these Γ functions, and obtain the Regge-pole structure [11] of the amplitude.

Presumably, other similar linear relations as Eq.(6) for high energy scatterings can be derived for Dp -brane with $p \geq 1$. However, in the following, in contrast to the common wisdom [15], we will show that, instead of the exponential fall-off of the form factors with Regge-pole structure, the high-energy scattering amplitudes of string scattered from D24-brane, i.e. Domain-wall, behave like *power-law* with Regge-pole structure. Moreover, we discover that the expected linear relations of high-energy string scattering amplitudes at each fixed mass level, Eq.(5) or Eq.(6), break down for the Domain-wall scatterings. We consider an incoming tachyon closed string state with momentum k_1 and an angle of incidence ϕ and an outgoing massive closed string state $(\alpha_{-1}^T)^{n-2q} (\alpha_{-2}^L)^q \otimes (\tilde{\alpha}_{-1}^T)^{n-2q'} (\tilde{\alpha}_{-2}^L)^{q'} |0\rangle$ with momentum k_2 and an angle of reflection θ . The kinematic setup is

$$e^P = \frac{1}{M} (-E, k_2 \cos \theta, -k_2 \sin \theta) = \frac{k_2}{M}, \quad (8)$$

$$e^L = \frac{1}{M} (-k_2, E \cos \theta, -E \sin \theta), \quad (9)$$

$$e^T = (0, \sin \theta, \cos \theta), \quad (10)$$

$$k_1 = (E, -k_1 \cos \phi, -k_1 \sin \phi), \quad (11)$$

$$k_2 = (-E, k_2 \cos \theta, -k_2 \sin \theta). \quad (12)$$

In the high energy limit, the angle of incidence ϕ is identified to the angle of reflection θ , and e^P approaches e^L , $k_1, k_2 \simeq E$. For the case of Domain-wall scattering $Diag D_{\mu\nu} = (-1, 1, -1)$, and we have

$$a_0 \equiv k_1 \cdot D \cdot k_1 \sim -2E^2 \sin^2 \phi - 2M_1^2 \cos^2 \phi + M_1^2, \quad (13)$$

$$\begin{aligned} b_0 &\equiv 2k_1 \cdot k_2 + 1 \\ &\sim 4E^2 \sin^2 \phi + 4M_1^2 \cos^2 \phi - (M_1^2 + M^2) + 1, \end{aligned} \quad (14)$$

The scattering amplitude can be calculated to be

$$\begin{aligned}
A_{D-Wall} = & (-1)^{q+q'} \int d^2 z_1 d^2 z_2 (z_1 - \bar{z}_1)^{k_1 \cdot D \cdot k_1} (z_2 - \bar{z}_2)^{k_2 \cdot D \cdot k_2} |z_1 - z_2|^{2k_1 \cdot k_2} |z_1 - \bar{z}_2|^{2k_1 \cdot D \cdot k_2} \\
& \cdot \left[\frac{ie^T \cdot k_1}{z_1 - z_2} + \frac{ie^T \cdot D \cdot k_1}{\bar{z}_1 - z_2} + \frac{ie^T \cdot D \cdot k_2}{\bar{z}_2 - z_2} \right]^{n-2q} \\
& \cdot \left[\frac{ie^T \cdot D \cdot k_1}{z_1 - \bar{z}_2} + \frac{ie^T \cdot k_1}{\bar{z}_1 - \bar{z}_2} + \frac{ie^T \cdot D \cdot k_2}{z_2 - \bar{z}_2} \right]^{n-2q'} \\
& \cdot \left[\frac{e^L \cdot k_1}{(z_1 - z_2)^2} + \frac{e^L \cdot D \cdot k_1}{(\bar{z}_1 - z_2)^2} + \frac{e^L \cdot D \cdot k_2}{(\bar{z}_2 - z_2)^2} \right]^q \left[\frac{e^L \cdot D \cdot k_1}{(z_1 - \bar{z}_2)^2} + \frac{e^L \cdot k_1}{(\bar{z}_1 - \bar{z}_2)^2} + \frac{e^L \cdot D \cdot k_2}{(z_2 - \bar{z}_2)^2} \right]^{q'}.
\end{aligned} \tag{15}$$

Set $z_1 = iy$ and $z_2 = i$ to fix the $SL(2, R)$ gauge, and include the Jacobian $d^2 z_1 d^2 z_2 \rightarrow 4(1-y^2)dy$, we have, for the $(0 \rightarrow 1)$ channel,

$$\begin{aligned}
& A_{D-Wall}^{(0 \rightarrow 1)} \\
& \simeq 4(2i)^{k_1 \cdot D \cdot k_1 + k_2 \cdot D \cdot k_2} \left(\frac{E \sin 2\phi}{2} \right)^{2n} \left(\frac{1}{2M \cos^2 \phi} \right)^{q+q'} \\
& \cdot \sum_{i=0}^{q+q'} \binom{q+q'}{i} 2^i \int_0^1 dy y^{k_2 \cdot D \cdot k_2} (1-y)^{2k_1 \cdot k_2 + 1} \\
& \cdot (1+y)^{2k_1 \cdot D \cdot k_2 + 1} \left[\frac{1+y}{1-y} \right]^{2n-(q+q')} \left(\frac{1}{1-y} \right)^i \\
& \simeq \left(\frac{E \sin 2\phi}{2} \right)^{2n} \left(\frac{1}{2M \cos^2 \phi} \right)^{q+q'} \\
& \cdot \sum_{i=0}^{q+q'} \binom{q+q'}{i} \cdot B\left(a_0 + 1, \frac{b+1}{2}\right) F_i
\end{aligned} \tag{16}$$

where

$$b = b_0 + n_b = b_0 - 2n + (q + q') - i, \tag{17}$$

$$F_i \equiv \left(1 + \sqrt{\left| \frac{b}{2a_0 + b} \right|} \right)^i + \left(1 - \sqrt{\left| \frac{b}{2a_0 + b} \right|} \right)^i \tag{18}$$

$$\simeq \left[(1 + 2C_i E \sin \phi)^i + (1 - 2C_i E \sin \phi)^i \right] \tag{19}$$

with

$$C_i \equiv \sqrt{\left| \frac{1}{M_1^2 - M^2 + 1 - 2n + (q + q') - i} \right|}. \tag{20}$$

F_i in Eq.(18) is the high energy limit of the generalized hypergeometric function ${}_3F_2\left(\frac{b+1}{2}, -\left[\frac{i}{2}\right], \frac{1}{2} - \left[\frac{i}{2}\right]; a_0 + \frac{b+3}{2}, \frac{1}{2}; 1\right)$ [17]. At this stage, it is crucial to note that

$$b \simeq b_0 \simeq -2a_0 \quad (21)$$

in the high energy limit for the Domain-wall scatterings. As a result, F_i reduces to the form of Eq.(19), and depends on the energy E . Thus in contrast to the generic Dp -brane scatterings with $p \geq 0$, which contain two independent kinematic variables, there is only one kinematic variable for the special case of Domain-wall scatterings. As we will see in the following calculation, this peculiar property will reduce the high energy beta function in Eq.(16) from exponential to power-law behavior and, simultaneously, breaks down the linear relations as we had in Eq.(6) for the D-particle scatterings. Another way to see this breakdown is by examining Eq.(5). One notes that the argument in the paragraph after Eq.(1) for scattering with two kinematic variables, which leads to Eq.(5), is no longer valid as there is only one kinematic variable for the Domain-wall scatterings. It thus becomes meaningless to study high-energy, fixed angle scattering process for the Domain-wall scatterings. Finally, the scattering amplitude for the $(0 \rightarrow 1)$ channel can be calculated to be (similar result can be obtained for the $(1 \rightarrow \infty)$ channel)

$$\begin{aligned} A_{D-Wall}^{(0 \rightarrow 1)} &\simeq \left(\frac{E \sin 2\phi}{2}\right)^{2n} \left(\frac{1}{2M \cos^2 \phi}\right)^{q+q'} B\left(a_0 + 1, \frac{b_0 + 1}{2}\right) \\ &\cdot \sum_{i=0}^{q+q'} \binom{q+q'}{i} \cdot \frac{(a_0)_0 \left(\frac{b_0}{2}\right)_{n_b/2}}{\left(a_0 + \frac{b_0}{2}\right)_{n_b/2}} (1 + 2C_i E \sin \phi)^i \end{aligned} \quad (22)$$

$$\begin{aligned} &\simeq \left(\frac{\cos \phi}{\sqrt{2}}\right)^{2n} \left(\frac{E \sin \phi}{M \sqrt{|M_1^2 - 2M^2 - 1|} \cos^2 \phi}\right)^{q+q'} \\ &\cdot \frac{\Gamma(a_0 + 1) \Gamma(\frac{b_0+1}{2})}{\Gamma(a_0 + \frac{b_0}{2} + \frac{3}{2})} \frac{1}{\left(\frac{M_1^2 - M^2 + 1}{2}\right)_{-n}} \end{aligned} \quad (23)$$

where $(\alpha)_n \equiv \frac{\Gamma(\alpha+n)}{\Gamma(\alpha)}$ for integer n . On the other hand, since the argument of $\Gamma(a_0 + 1)$ in

Eq.(23) is negative in the high-energy limit, we have, by using Eq.(7) and Eq.(21),

$$\begin{aligned} \frac{\Gamma(a_0 + 1)\Gamma(\frac{b_0+1}{2})}{\Gamma(a_0 + \frac{b_0}{2} + \frac{3}{2})} &\simeq \frac{\pi}{\sin(\pi a_0)} \frac{\Gamma(\frac{b_0+1}{2})}{\Gamma(-a_0)\Gamma(\frac{M_1^2 - M^2 + 1}{2})} \\ &\sim \frac{1}{\sin(\pi a_0)} \frac{1}{(E \sin \phi)^{2(n-1)}}. \end{aligned} \quad (24)$$

Note that the $\sin(\pi a_0)$ factor in the denominator of Eq.(24) gives the Regge-pole structure, and the energy dependence $E^{-2(n-1)}$ gives the power-law behavior in the high energy limit. As a result, the scattering amplitude for the Domain-wall in Eq.(23) behaves like power-law with the Regge-pole structure. The crucial differences between the Domain-wall scatterings in Eq.(23) and the D-particle scatterings (or any other Dp -brane scatterings except Domain-wall and D-instanton scatterings) in Eq.(6) is the kinematic relation Eq.(21). For the case of D-particle scatterings [17], the corresponding factors for both F_i in Eq.(18) and the fraction in Eq.(22) are independent of energy in the high energy limit, and, as a result, the amplitudes contain no $q + q'$ dependent energy power factor. So one gets the high energy linear relations for the D-particle scattering amplitudes. On the contrary, for the case of Domain-wall scatterings, both F_i in Eq.(19) and the fraction in Eq.(22) depend on energy due to the condition Eq.(21). The summation in Eq.(22) is then dominated by the term $i = q + q'$, and the whole scattering amplitude Eq.(23) contains a $q + q'$ dependent energy power factor. As a result, the usual linear relations for the high energy scattering amplitudes break down for the Domain-wall scatterings. It is crucial to note that the mechanism, Eq.(21), to drive the exponential fall-off form factor of the D-particle scatterings to the power-law one of the Domain-wall scatterings is exactly the same as the mechanism to break down the expected linear relations for the domain-wall scatterings in the high energy limit. In conclusion, this result gives a strong evidence that the existence of the infinite linear relations, or stringy symmetries, of high-energy string scattering amplitudes is responsible for the softer, exponential fall-off high-energy string scatterings than the power-law field theory scatterings.

Another interesting case of D-brane scatterings is the massless form factor of scatterings of D-instanton [15]

$$\frac{\Gamma(s)\Gamma(t)}{\Gamma(s+t+1)} \rightarrow \frac{1}{st}, \text{ as } s \rightarrow 0, \quad (25)$$

which contains no Regge-pole structure. In Eq.(25), s, t are the Mandelstam variables. Eq.(25) can be easily generalized to the scatterings of arbitrary massive string states in the

high energy limit. To compare the D-instanton scatterings with the Domain-wall scatterings in Eq.(24), one notes that in both cases there is only one kinematic variable and, as a result, behave as power-law at high energies [19]. On the other hand, since t is large negative in the high energy limit [11], the application of Eq.(7) to Eq.(25) produces no $\sin(\pi a_0)$ factor in contrast to the Domain-wall scatterings. So there is no Regge-pole structure for the D-instanton scatterings. We conclude that the very condition of Eq.(21) makes Domain-wall scatterings an unique example of a hybrid of string and field theory scatterings.

This work is supported in part by the National Science Council, 50 billions project of MOE and National Center for Theoretical Science, Taiwan, R.O.C.

-
- [1] D. J. Gross and P. F. Mende, Phys. Lett. B **197**, 129 (1987); Nucl. Phys. B **303**, 407 (1988).
 - [2] D. J. Gross, Phys. Rev. Lett. **60**, 1229 (1988); Phil. Trans. R. Soc. Lond. A329, 401 (1989).
 - [3] D. J. Gross and J. L. Manes, Nucl. Phys. B **326**, 73 (1989). See section 6 for details.
 - [4] C. T. Chan and J. C. Lee, Phys. Lett. B **611**, 193 (2005). J. C. Lee, [arXiv:hep-th/0303012].
 - [5] C. T. Chan and J. C. Lee, Nucl. Phys. B **690**, 3 (2004).
 - [6] C. T. Chan, P. M. Ho and J. C. Lee, Nucl. Phys. B **708**, 99 (2005).
 - [7] C. T. Chan, P. M. Ho, J. C. Lee, S. Teraguchi and Y. Yang, Nucl. Phys. B **725**, 352 (2005).
 - [8] C. T. Chan, P. M. Ho, J. C. Lee, S. Teraguchi and Y. Yang, Phys. Rev. Lett. **96** (2006) 171601.
 - [9] C. T. Chan, P. M. Ho, J. C. Lee, S. Teraguchi and Y. Yang, Nucl. Phys. B **749**, 266 (2006).
 - [10] C. T. Chan, J. C. Lee and Y. Yang, Nucl. Phys. B **738**, 93 (2006).
 - [11] C. T. Chan, J. C. Lee and Y. Yang, Nucl. Phys. B **749**, 280 (2006).
 - [12] J. C. Lee, Phys. Lett. B **241**, 336 (1990); Phys. Rev. Lett. **64**, 1636 (1990). J. C. Lee and B. Ovrut, Nucl. Phys. B **336**, 222 (1990); J.C.Lee, Phys. Lett. B **326**, 79 (1994).
 - [13] T. D. Chung and J. C. Lee, Phys. Lett. B **350**, 22 (1995). Z. Phys. C **75**, 555 (1997). J. C. Lee, Eur. Phys. J. C **1**, 739 (1998).
 - [14] H. C. Kao and J. C. Lee, Phys. Rev. D **67**, 086003 (2003). . C. T. Chan, J. C. Lee and Y. Yang, Phys. Rev. D **71**, 086005 (2005)
 - [15] For a review, see A. Hashimoto and I.R. Klebanov, "Scattering of strings from D-branes" hep-th/9611214.
 - [16] M.R. Garousi and R.C. Myers, "Superstring scattering from D-Branes" Nucl. Phys. B475

- (1996) 193, hep-th/9603194.
- [17] C. T. Chan, J. C. Lee and Y. Yang, " Scatterings of massive string states from D-brane and their linear relations at high energies", Nucl.Phys.B**764**, 1 (2007).
- [18] M. B. Green, J. H. Schwarz and E. Witten, "Superstring Theory. Vol. 1," Cambridge University Press 1987.
- [19] It is worth noting that the amplitude for two massless D-brane fluctuations to produce a massless closed string contains only one kinematic invariant [15]. However, the high energy limit of this amplitude behaves as exponential fall-off instead of power-law. See J.C.Lee and Y. Yang, Phys.Lett. B646 (2007) 120.